

About calculation of the Hankel transform using preliminary wavelet transform

E.B. Postnikov

Theoretical Physics Department, Kursk State Pedagogical University,
Radischsheva st. 33, Kursk, Russia, 305000
e-mail: postnicov@mail.ru

MSC-class: 44A15, 65R10

Abstract

The purpose of this paper is to present an algorithm for evaluating Hankel transform of the null and the first kind. The result is the exact analytical representation as the series of the Bessel and Struve functions multiplied to the wavelet coefficients of the input function. Numerical evaluation of the test function with known analytical Hankel transform illustrates proposed algorithm.

The Hankel transform is a very useful instrument in a wide range of physical problems which have an axial symmetry [1]. The influence of the Laplasian on a function in a cylindrical coordinates is equal to the product of the parameter of the transformation squared and the transform of the function:

$$\begin{cases} \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) f(r) \leftrightarrow -p^2 F_0(p) \\ \left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{1}{r^2} \right) f(r) \leftrightarrow -p^2 F_1(p) \end{cases} \quad (1)$$

The Hankel transform of the null ($n = 0$) and the first ($n = 1$) kind are represented as

$$\begin{aligned} F_n(p) &= \int_0^\infty f(r) J_n(pr) r dr, \\ f_n(p) &= \int_0^\infty F(p) J_n(pr) p dp. \end{aligned} \quad (2)$$

Besides that integrals like (2) are connected with the problems of geophysics and cosmology, for example [3, 4].

But practical calculation of direct and inverse Hankel transform is connected with two problems. The first problem is based on the fact that not every transform in the real physical situation has analytical expression for result of inverse Hankel transform. The second one is determination of functions as a set of their values for numerical calculations. Large bibliography on those issues can be found in [5]. The classical trapezoidal rule, Cotes rule and other rules connected with replacement of integrand by sequence of polynoms have high accuracy if integrand is a smooth function. But $f(r)J_n(pr)r$ (or $F_p(p)J_n(pr)p$) is a quick oscillating function if r (or p) is large. There are two general methods of the effective calculation in this area. The first is the Fast Hankel transform [6]. The specification of that method is transforming the function to the logarithmical space and fast Fourier transform in that space. This method needs a smoothing of the function in log-space. The second method is based on the separation of the integrand into product of slowly

varying component and a rapidly oscillating Bessel function [7]. But it needs the smoothness of the slow component for its approximation by low-order polynoms.

The goal of this article is to apply wavelet transform with Haar bases to (2).

The both direct and inverse transforms (2) are symmetric. Let us consider only one of them, for example, direct transform. Let's denote $f(r)r$ as $g(r)$. Then Hankel transform is

$$F_{0,1}(p) = \int_0^\infty g(r) J_{0,1}(pr) dr \quad (3)$$

The expansion $g(r) \in L^2(R)$ into wavelet series with the Haar bases is [8] :

$$g(r) = \sum_{k=0}^\infty c_{0k} \varphi_k(r) + \sum_{j=0}^\infty \sum_{k=0}^\infty d_{jk} \psi_{jk}(r), \quad (4)$$

$$\varphi_{0k}(r) = \varphi^H(r - k), \psi_{jk} = 2^{j/2} \psi^H(2^j r - k),$$

$$\varphi^H(t) = \begin{cases} 1, & t \in (0, 1) \\ 0, & t \notin (0, 1) \end{cases}, \psi^H(t) = \begin{cases} 1, & t \in (0, 1/2) \\ -1, & t \in (1/2, 0) \\ 0, & t \notin (0, 1) \end{cases}.$$

After substitution (4) into (3) one has

$$F_{0,1}(p) = \sum_{k=0}^\infty c_{0k} \int_0^\infty \varphi_k(r) J_{0,1}(pr) dr + \sum_{j=0}^\infty \sum_{k=0}^\infty d_{jk} \int_0^\infty \psi_{jk}(x) J_{0,1}(pr) dr.$$

Making use integrals from [2] we have as a result

$$\begin{aligned} F_0(p) &= \sum_{k \in \mathbb{Z}} c_{0k} [(k+1)J_0(p(k+1)) - kJ_0(pk) + \\ &\frac{\pi}{2} [(k+1)D(p(k+1)) - kD(pk)] + \\ &\sum_{j=0}^\infty \sum_{k \in \mathbb{Z}} d_{jk} [2(k+\frac{1}{2})J_0(p(k+\frac{1}{2})) - (k+1)J_0(p(k+1)) - kJ_0(pk) - \\ &\frac{\pi}{2} [2(k+\frac{1}{2})D(p(k+\frac{1}{2})) - (k+1)D(p(k+1)) - kD(pk)] \end{aligned} \quad (5)$$

$$\begin{aligned} F_1(p) &= \frac{1}{p} \left\{ \sum_{k \in \mathbb{Z}} c_{0k} [J_0(pk) - J_0(p(k+1))] + \right. \\ &\left. \sum_{j=0}^\infty \sum_{k \in \mathbb{Z}} d_{jk} [2J_0(p(k+\frac{1}{2})2^{-j}) - J_0(p(k+1)2^{-j}) - J_0(pk2^{-j})] \right\}, \quad (6) \end{aligned}$$

where $D(\xi) = H_0(\xi)J_1(\xi) - H_1(\xi)J_0(\xi)$ and $H_{0,1}$ is Struve function of the null and the first kind.

The most sufficient result is that equations (5) and (6) are exact. They can be used in any analytical expressions. Especially it's useful for Hankel transform of the first kind because (6) contains only a combination of Bessel functions, and one can use such their properties as orthogonality etc. The coefficients c_{0k} means average value of $g(r)$ at the range $[k, k+1]$:

$$c_{0k} = \int_k^{k+1} g(r) dr$$




Figure 1: Original function (left) and Hankel transform (right)

The detail coefficients are

$$d_{jk} = 2^{j/2} \left\{ \int_{2^{-j}k}^{2^{-j}(k+\frac{1}{2})} g(r)dr - \int_{2^{-j}(k+\frac{1}{2})}^{2^{-j}(k+1)} g(r)dr \right\}.$$

The formulas (5) and (6) allow us to get full analytical solution if integrals above have close form solution. In the opposite case the solution must be numerical but this method provides an effective algorithm for that. It's obvious that d_{jk} decrease very quickly if $g(r)$ is a smooth function. One can practically use $d_{jk} > \varepsilon$, where ε is small. But if $g(r)$ has steps, sharp vertices or discontinues then the detail coefficients concentrate around these points and one can appropriate they are equal to the zero in other areas.

Let us consider for example a function with known analytical Hankel transform

$$\int_0^{\infty} e^{-a^2 r^2} r J_1(pr) r dr = \frac{p}{4a^4} e^{-p^2/4a^2}. \quad (7)$$

The approximation and detail coefficients may be calculated analytically in closed form

$$\begin{aligned} c_{0k} &= \left. \frac{\sqrt{\pi} \operatorname{erf}(r) - 2a r e^{-a^2 r^2}}{4a^3} \right|_k^{(k+1)} \\ d_{jk} &= 2^{j/2} \left. \frac{\sqrt{\pi} \operatorname{erf}(r) - 2a r e^{-a^2 r^2}}{4a^3} \right|_{2^{-j}k}^{2^{-j}(k+\frac{1}{2})} - \\ &\quad 2^{j/2} \left. \frac{\sqrt{\pi} \operatorname{erf}(r) - 2a r e^{-a^2 r^2}}{4a^3} \right|_{2^{-j}(k+\frac{1}{2})}^{2^{-j}(k+1)} \end{aligned} \quad (8)$$

Thus (6) with the coefficients (8) is the exact representation of the Hankel transform. Let us consider the approximate solution. Suppose the function is known (7) only in the segment $[0, h]$. Then there is the series instead of (4):

$$g(r) = c_0 \varphi_0(r) + \sum_{j=0}^J \sum_{k=0}^{2^j-1} d_{jk} \psi_{jk}(r). \quad (9)$$

If $J \rightarrow \infty$ then (9) is exact for this truncated function. But practically we only use several first levels. For example we can see the original function (the replacement r to $x = r/h$ is used) (7) and the transform at the Fig.1. One can see that exact transform (solid line) and the transform at level $J = 3$ (dotted line) coincide at




Figure 2: Transform's error

this figure. The absolute errors between the exact transform and the approximate transform at the levels $J = 2$ (solid line), $J = 3$ (dashed line), and $J = 4$ (dotted line) are represented at the Fig.2 (left). It's obvious that the error is small in comparison with the values of the $F_1(p)$. The absolute error at the level $J = 3$ in a wide range of p is plotted in the right side of the Fig.2. One can see that this error has quasi-periodic oscillations because the function is truncated. But they decrease with the growth of p (and J) when oscillations in classical Fast Hankel Transform [4] increase.

References

- [1] J. Mathews, R.L. Walker Mathematical methods of physics. New-York- Amsterdam, W.A. Benjamin, Inc, 1964
- [2] M. Abramowitz and I. A. Stegun (eds.), Handbook of mathematical functions with formulas, graphs and mathematical tables, National Bureau of Standards Applied Mathematics Series, vol. 55, U. S. Government Printing Office, Washington, D. C., 1964.
- [3] J. Zhao, W.W.M. Dai, S. Kapur, D.E. Long Efficient Three-Dimensional Extraction Based on Statuic and Full-Wave Layered Green's Function. Proceedings of the 35 Conference on Design Automation, Moscon center, San Francisco, USA, June 15-19, 1998. ACM Press, 1998, pp. 224-229
- [4] Hamilton A.J.S. Uncorrelated modes of the Nonlinear Power Spectrum. Mon. Not. Roy. Astron. Soc. 2000, Iss. 312, pp. 257-284.
- [5] D. W. Lozier and F. W. J. Olver, Numerical evaluation of special functions, Mathematics of Computation 1943-1993: A Half-Century of Computational Mathematics (W. Gautschi, ed.), Proceedings of Symposia in Applied Mathematics, vol. 48, American Mathematical Society, Providence, Rhode Island 02940, 1994, pp. 79-125. See also <http://math.nist.gov/nesf/>.
- [6] A. E. Siegman, Quasi fast Hankel transform, Optics Lett. 1 (1977), pp. 13-15.
- [7] R. Barakat and E. Parshall, Numerical evaluation of the zero-order Hankel transform using Filon quadrature philosophy, Appl. Math. Lett. 9 (1996), no. 5, pp. 21-26.
- [8] C.K. Chui An introduction to Wavelets. Academic Press, 1992.